

# Neuron

## Purification and Characterization of Progenitor and Mature Human Astrocytes Reveals Transcriptional and Functional Differences with Mouse

### Highlights

- We develop the first method to acutely purify fetal and adult human astrocytes
- We obtain transcriptome profiles of human neurons, glia, and vascular cells
- While similar, human and mouse astrocytes have unique genomic and functional traits
- Human astrocytes exist in at least two distinct developmental stages

### Authors

Ye Zhang, Steven A. Sloan, Laura E. Clarke, ..., Gerald A. Grant, Melanie G. Hayden Gephart, Ben A. Barres

### Correspondence

zhangye@stanford.edu

### In Brief

Zhang et al. developed a method to acutely purify healthy and diseased human astrocytes and to culture them in serum-free conditions. They obtained transcriptome profiles of purified human CNS cell types and discovered two distinct stages of human astrocyte development.

### Accession Numbers

GSE73721

# Purification and Characterization of Progenitor and Mature Human Astrocytes Reveals Transcriptional and Functional Differences with Mouse

Ye Zhang,<sup>1,7,\*</sup> Steven A. Sloan,<sup>1,7</sup> Laura E. Clarke,<sup>1</sup> Christine Caneda,<sup>1</sup> Colton A. Plaza,<sup>1</sup> Paul D. Blumenthal,<sup>2</sup> Hannes Vogel,<sup>3</sup> Gary K. Steinberg,<sup>4</sup> Michael S.B. Edwards,<sup>4</sup> Gordon Li,<sup>4</sup> John A. Duncan III,<sup>5</sup> Samuel H. Cheshier,<sup>4</sup> Lawrence M. Shuer,<sup>4</sup> Edward F. Chang,<sup>6</sup> Gerald A. Grant,<sup>4</sup> Melanie G. Hayden Gephart,<sup>4</sup> and Ben A. Barres<sup>1</sup>

<sup>1</sup>Department of Neurobiology, Stanford University School of Medicine, Stanford, CA 94305, USA

<sup>2</sup>Department of Obstetrics and Gynecology, Stanford University Medical Center, Stanford, CA 94305, USA

<sup>3</sup>Department of Pathology, Stanford University Medical Center, Stanford, CA 94305, USA

<sup>4</sup>Department of Neurosurgery, Lucile Packard Children's Hospital and Stanford University Medical Center, Stanford, CA 94305, USA

<sup>5</sup>Department of Pediatric Neurosciences, Kaiser Permanente Santa Clara Medical Center, Santa Clara, CA 95051, USA

<sup>6</sup>UCSF Epilepsy Center, University of California, San Francisco, 400 Parnassus Avenue, San Francisco, CA 94143, USA

<sup>7</sup>Co-first author

\*Correspondence: [zhangye@stanford.edu](mailto:zhangye@stanford.edu)

<http://dx.doi.org/10.1016/j.neuron.2015.11.013>

## SUMMARY


The functional and molecular similarities and distinctions between human and murine astrocytes are poorly understood. Here, we report the development of an immunopanning method to acutely purify astrocytes from fetal, juvenile, and adult human brains and to maintain these cells in serum-free cultures. We found that human astrocytes have abilities similar to those of murine astrocytes in promoting neuronal survival, inducing functional synapse formation, and engulfing synaptosomes. In contrast to existing observations in mice, we found that mature human astrocytes respond robustly to glutamate. Next, we performed RNA sequencing of healthy human astrocytes along with astrocytes from epileptic and tumor foci and compared these to human neurons, oligodendrocytes, microglia, and endothelial cells (available at <http://www.brainrnaseq.org>). With these profiles, we identified novel human-specific astrocyte genes and discovered a transcriptome-wide transformation between astrocyte precursor cells and mature post-mitotic astrocytes. These data represent some of the first cell-type-specific molecular profiles of the healthy and diseased human brain.

## INTRODUCTION

Constituting about 40% of all cells in the human brain, astrocytes have long been classified as mere passive support cells. Recent work, however, has demonstrated that murine astrocytes play many active roles and are critical for the development and function of the CNS (Allen and Barres, 2009; Chung et al., 2015; Clarke and Barres, 2013; Hamilton and Attwell, 2010; Haydon

and Nedergaard, 2014; Khakh and McCarthy, 2015; Molofsky et al., 2012). For example, purified neurons in culture have little ability to form functional synapses, which are powerfully promoted by astrocytes (Allen et al., 2012; Christopherson et al., 2005; Kucukdereli et al., 2011; Ullian et al., 2001). Murine astrocytes are not only important for the formation and function of synapses, but are also essential for the phagocytic elimination of synapses and the refinement of developing neural circuits (Chung et al., 2013). Since synapse formation, function, and elimination are key processes occurring during learning and memory, astrocytes are postulated to be an indispensable component in CNS plasticity. Additionally, murine astrocytes are required for neurotransmitter recycling (Rothstein et al., 1996), extracellular potassium homeostasis (Kuffler et al., 1966), regulation of blood flow (Attwell et al., 2010; Mulligan and MacVicar, 2004; Zonta et al., 2003), and providing energy substrates for neurons (Pellerin and Magistretti, 1994). Considering their central role in CNS physiology, it is not surprising that astrocyte dysfunction has been demonstrated or implicated in nearly all neurological disorders (Molofsky et al., 2012). Currently, the extent of our understanding of astrocyte physiology in health and disease is almost entirely restricted to observations in murine models, but how primary human and murine astrocytes compare at molecular and functional levels remains largely unknown.

Observational studies from postmortem human tissues have revealed that human astrocytes are much larger and more complex than their murine counterparts (Oberheim et al., 2006, 2009). More recently, transplantation of human glial progenitors into mouse brains has been shown to improve learning and memory (Han et al., 2013). These observations raise questions about how murine astrocyte physiology and function might extend to humans. Are there distinct properties of human astrocytes that contribute to human intelligence and cognition? Can we extend findings about astrocyte pathology in neurological disorders from mouse and rat studies to develop effective therapeutic approaches in humans?



**Figure 1. Acute Purification of Fetal and Mature Human Astrocytes**

(A) Juvenile and adult (8–63 years old; yo) patient temporal lobe cortex tissue and fetal (17–20 gw old) brain tissue were first dissociated into single-cell suspensions.

(legend continued on next page)

A major hurdle in addressing these issues is the lack of a method to acutely purify human astrocytes and culture them in chemically defined conditions. Current purification methods for human astrocytes require culturing dissociated nervous tissue in serum for days to weeks (McCarthy and de Vellis, 1980). In vivo, however, quiescent astrocytes do not contact serum except upon injury and blood-brain-barrier breakdown, and in vitro exposure to serum has been shown to induce irreversible reactive changes in astrocytes (Foo et al., 2011; Zamanian et al., 2012). Moreover, since serum-selection methods require a group of proliferating astrocyte progenitors, these protocols do not work efficiently to purify mature astrocytes from adult human brains. Because of these limitations, the transcriptome profile of mature resting human astrocytes has remained difficult to ascertain.

The method we present here is an immunopanning-based technique that utilizes an antibody targeted against HepaCAM (or GlialCAM), a surface protein expressed by human astrocytes, to generate purified (>95%) cultures of primary human astrocytes. With this method, we acutely purified astrocytes from over 20 fetal, juvenile, and adult human subjects and then performed transcriptome profiling and functional studies to compare developmental and interspecies differences. Additionally, we purified each of the major CNS cell types and generated a user-friendly searchable online database to enable dissemination of human cell-type-specific gene expression data to the greater neuroscience community (<http://www.brainrbaseq.org>).

## RESULTS

### Acute Purification and Serum-free Culture of Fetal and Mature Human Astrocytes

We obtained juvenile (8–18 years old) and adult (21–63 years old) human brain tissue samples from patients undergoing neurological surgeries, with informed consent under a protocol approved by the Stanford University Institutional Review Board. The samples used in this study (except for the data shown in Figures 5 and 6 for epilepsy and glioblastoma) were small pieces of healthy temporal lobe cortices that were resected to gain access to deeper epileptic hippocampi (Figure 1A). In each of these cases, the temporal lobe cortex that was removed was considered normal based upon MRI, electroencephalogram (EEG) studies, and pathological examination. Detailed information on patient diagnoses is listed in Table S1. We worked closely with the surgeons to minimize the time from tissue resection to the beginning of the purification protocol (within 1 hr). Furthermore, we generated all gene expression data by collecting RNA from cells immediately after purification to best represent their in vivo molecular profiles. In addition to surgical samples, we also obtained fetal brain tissue from elective pregnancy terminations at 17–20 gestational weeks under a protocol approved by the Stanford University Institutional Review Board. For fetal cases, we began purification within 5 hr after the procedures.


To develop a method for the purification of astrocytes, we first sought to identify an astrocyte-specific antibody that we could use to immunopan human cortical tissue. We mined our existing mouse astrocyte RNA sequencing (RNA-seq) datasets (Zhang et al., 2014) for potential surface markers that were enriched in astrocytes, not expressed by radial glial cells (a major cellular constituent of fetal human brain), and where antibodies to the human antigen already existed. After screening numerous candidates, we settled on HepaCAM, a cell-adhesion glycoprotein that is enriched in astrocytes. The immunopanning protocol involves passing a single-cell suspension of dissociated tissue over a series of Petri dishes coated with antibodies directed against cell-type-specific antigens (Figures 1A and 1B). For juvenile and adult samples, our immunopanning protocol consisted of anti-CD45 antibodies to bind microglia (and macrophages), anti-GalC hybridoma to bind oligodendrocytes and myelin debris, anti-O4 hybridoma to bind oligodendrocyte precursor cells (OPCs), anti-Thy1 antibody to bind neurons, anti-HepaCAM antibody to bind astrocytes, and *Banisteria simplicifolia* lectin 1 (BSL-1) to bind endothelial cells. We used a shortened antibody-binding procedure for fetal samples, because oligodendrocytes and myelin are not yet generated at 17–20 gestational weeks (Experimental Procedures; Supplemental Information). After cells were bound to an antibody-coated dish, we washed away loosely bound contaminating cells and then either detached cells via trypsin digestion for culture or scraped cells directly off the dish to extract RNA for RNA-seq.

We routinely obtained over 95% pure mature human astrocytes as determined by glial fibrillary acidic protein (GFAP) immunofluorescence (Figure 1D) and further validated these immunohistochemical purity estimates by RNA-seq (Figure 4; also discussed later). We refer to astrocytes purified from juvenile and adult brain tissue as “mature” astrocytes to distinguish them from fetal astrocytes. Mature human astrocytes exhibited process-bearing morphologies (Figure 1F; Figure S1) and remained healthy for 3 weeks or longer in vitro when cultured in serum-free defined media (Supplemental Information). Interestingly, while fetal human astrocytes proliferated robustly in vitro, mature astrocytes did not divide. We performed immunostaining of fetal and mature human astrocytes with an antibody against a mitotic cell marker, Ki67. While  $14.9\% \pm 1.6\%$  of fetal human astrocytes were actively proliferating, only  $0.05\% \pm 0.1\%$  of cells in mature human astrocyte cultures were positive for Ki67. Similarly, we performed an EdU (5-ethynyl-2'-deoxyuridine) incorporation assay and detected EdU labeling in  $15.0\% \pm 2.3\%$  of fetal human astrocytes and only  $0.4\% \pm 0.5\%$  of cells in mature human astrocyte cultures (Figure S2). This was further confirmed by mRNA expression of two mitotic markers, TOP2A and MKI67, which were exclusively present in fetal astrocytes. (For TOP2A: FPKM [fragments per kilobase of transcript sequence per million mapped fragments] =  $38.2 \pm 6.2$  in fetal astrocytes and  $0.1 \pm 0.0$  in mature astrocytes; for MKI67: FPKM =  $34.2 \pm 3.9$  in fetal astrocytes and  $0.1 \pm 0.0$  in

(B) Schematics of immunopanning purification of cell types from human brain samples.

(C and D) Unpurified brain cells (left), Thy1-purified neurons (middle), and HepaCAM-purified astrocytes (right) from fetal (C) and mature (D) brains stained at 7 d.i.v. for neurons (TuJ1, red), astrocytes (GFAP, green), and nuclei of all cells (DAPI, blue). Scale bars, 100  $\mu$ m.

(E and F) Cultured human fetal (E) and mature (F) astrocytes grown in culture for 7 days and stained with GFAP. Scale bars, 100  $\mu$ m; scale bars in insets, 50  $\mu$ m.



**Figure 2. Functional Characterization of Human Astrocytes**

(A) Schematics of co-culture experiments. 17–20 gw fetal human astrocytes and neurons were purified by immunopanning, grown in the same wells separated by porous inserts.

(B) Calcein stain of live neurons (green) and ethidium homodimer stain of dead neurons (red) in the presence and absence of astrocytes. Scale bars, 100  $\mu$ m.

(legend continued on next page)



mature astrocytes. Data represent average  $\pm$  SEM, unless otherwise noted. For fetal astrocytes,  $n = 6$  patients; for mature astrocytes,  $n = 12$  patients.) Therefore, human astrocytes exist in two distinct developmental stages: a fetal, highly proliferative astrocyte precursor cell (APC) stage and a mature, post-mitotic stage. Due to their proliferative nature, acutely purified human APCs can be frozen, stored, and defrosted at a later time. Frozen and defrosted APCs remain healthy and continue to proliferate in culture at similar rates (Figure S2), so experiments using immunopanned human APCs are not time-locked to tissue availability.

### Human Astrocytes Promote Neuron Survival

Our purification technique allowed us to investigate the functional capabilities of primary human astrocytes *in vitro*. First, we tested whether human astrocytes could promote neuronal survival, a hallmark feature of mouse and rat astrocytes (Banker, 1980). We immunopanned human neurons and astrocytes (Figures 1A and 1B) and grew them together in the same cell-culture wells on separate layers. The inserts separating the two cell types allowed secreted molecules to freely diffuse while preventing direct astrocyte-to-neuron contact (Figure 2A). After 7 days of co-culture, we stained live neurons with calcein-AM and dead neurons with ethidium homodimer-1 (Figure 2B). We found that human astrocytes strongly promoted neuronal survival in a dose-dependent manner (neuronal survival:  $2.2\% \pm 0.9\%$  without astrocytes,  $88.5\% \pm 2.4\%$  with 200,000 astrocytes;  $n = 8$ ) (Figure 2C). Thus, human astrocytes share the ability of murine astrocytes to strongly promote neuronal survival *in vitro*.

### Human Astrocytes Promote Synapse Formation

To assess whether human astrocytes, like rat astrocytes, promote synapse formation, we co-cultured human astrocytes with rat retinal ganglion cells (RGCs), using inserts to separate the two cell types. We chose rat RGCs because our group previously developed a culture medium that supports RGC survival in the absence of astrocytes (Figures S3C and S3D) (Meyer-Franke et al., 1995; Winzeler and Wang, 2013). After co-culturing human astrocytes with RGCs for 14–17 days, we performed double immunostaining for the presynaptic marker Bassoon and the postsynaptic marker Homer to quantify colocalized synaptic puncta.

At the immunohistological level, we found that human astrocytes increased the number of synapses 3- to 4-fold, as we have previously reported for rat astrocytes. This was true for astrocytes of all developmental ages, including fetal, juvenile, and adult time points (Figure 2H). (Normalized synapse number:  $1.0 \pm 0.2$  without astrocytes,  $3.8 \pm 1.2$  with fetal astrocytes,  $3.1 \pm 0.3$  with juvenile astrocytes, and  $2.5 \pm 0.5$  with adult astrocytes.  $n = 10$ –20 images per patient. Each image contains, on average, two cells. Fetal: 1 patient; age: 18 gestational weeks [gw] old. Juvenile: 3 patients, ages: 8, 13, and 18 years old. Adult: 3 patients; ages: 26, 41, and 47 years old.) In addition to increasing the number of synapses, we also found that the area of colocalization of Homer/Bassoon positive puncta was significantly increased when human astrocytes were present. (Average colocalization pixel overlap =  $16.2 \pm 0.6$  without astrocytes and  $20.8 \pm 0.5$  with astrocytes.  $N_s = 1,065$  and  $3,212$  synapses in the absence and presence, respectively, of astrocytes. Data were obtained using astrocytes from 7 patients; ages: 18 gw and 8, 13, 18, 26, 41, and 47 years old.)

Next, to examine whether the synapses induced by human astrocytes were functional and to further validate the synaptogenic effects of human astrocytes, we performed patch-clamp recordings of neurons cultured with or without astrocyte feeder layers. Consistent with our immunohistochemical observations, we found that human astrocytes significantly increased the amplitude of miniature excitatory postsynaptic currents (mEPSCs) (rightward shift in the cumulative probability plot of mEPSC amplitude,  $p = 0.0001$ , Kolmogorov-Smirnov test; mean amplitude of RGCs alone:  $21.7 \pm 0.4$  pA; RGCs with human astrocytes:  $23.7 \pm 0.3$  pA;  $n_s = 488$  and  $2,837$  mEPSCs from 16 and 18 cells, for no astrocytes and with astrocytes, respectively; Figure 2J) and frequency of mEPSCs (RGCs alone:  $0.4 \pm 0.1$  Hz; RGCs with human astrocytes:  $1.9 \pm 0.4$  Hz,  $p = 0.004$ .  $n_s = 15$  and  $23$  cells for no astrocytes and with astrocytes, respectively; Figures 2I and 2K). Taken together, these results demonstrate that human astrocytes strongly induce the formation and function of excitatory synapses.

### Human Astrocytes Engulf Synaptosomes

Recent studies have found that mouse astrocytes are involved in eliminating extra synapses to help refine neural circuits (Chung

(C) Quantification of survival rate.  $N = 7$ –8 images per condition.  $**p < 0.01$ .  $***p < 0.001$ , two-tailed unpaired Student's *t* test.

(D) Human astrocytes engulf synaptosomes *in vitro*. FACS plot of human fetal astrocytes incubated without synaptosomes, with synaptosomes, or with synaptosomes and 5% serum.

(E) Percentages of synaptosome-positive astrocytes.  $*p = 0.05$ , one-tailed Wilcoxon rank-sum test.  $n = 3$  patients.

(F) Confocal image of a human fetal astrocyte stained with Vibrant CFDA (cyan) engulfing PhrodoRed-labeled synaptosomes (magenta) Scale bar, 20  $\mu$ m.

(G) Retinal ganglion cells form more synapses in the presence of human astrocytes. Cyan indicates immunofluorescence of post-synaptic marker, Homer. Magenta indicates immunofluorescence of pre-synaptic marker, Bassoon. Scale bar, 10  $\mu$ m.

(H) The number of synapses (Homer/Bassoon double-positive puncta) in the presence and absence of human astrocytes.  $N = 10$ –20 images per patient. Each image contains, on average, two cells. Fetal: 1 patient; age, 18 gw. Juvenile: 3 patients; ages, 8, 13, and 18 years old. Adult: 3 patients; ages: 26, 41, and 47 years old.  $**p < 0.01$ , two-tailed unpaired Student's *t* test.

(I) Representative traces of whole-cell patch-clamp recordings from RGCs cultured with or without feeder layers of human astrocytes in the presence of TTX. Few mEPSCs were observed without feeder layers of human astrocytes.

(J) Human astrocytes significantly increased the amplitude of mEPSCs ( $p = 0.0001$ , Kolmogorov-Smirnov test;  $n_s = 488$  and  $2,837$  mEPSCs from 16 and 18 cells, respectively, for no astrocytes and with astrocytes respectively).

(K) The frequency of mEPSCs was significantly increased in the presence of human astrocytes (two-tailed Wilcoxon rank-sum test.  $p = 0.004$ .  $n = 16$ –18 neurons per condition).  $**p < 0.01$ , two-tailed unpaired Student's *t* test.

Data represent mean  $\pm$  SEM unless otherwise noted.

et al., 2013). To address whether human astrocytes share similar capabilities, we performed an in vitro synaptosome engulfment assay. We purified synaptosomes and labeled them with a fluorescent dye, PhrodoRed. PhrodoRed is non-fluorescent at neutral pH and fluoresces brightly in acidic environments. Therefore, it allowed us to detect engulfed synaptosomes that had been trafficked to lysosomes with minimal background fluorescence from sticky synaptosomes attached to the astrocyte surface. To assess the amount of phagocytosis in human astrocytes, we fed PhrodoRed-labeled synaptosomes to purified astrocytes in the presence of either astrocyte-conditioned medium (ACM) or serum. Although serum contains high concentrations of bridging molecules that are required for phagocytosis, it is also known to induce reactive astrogliosis. Therefore, we wondered whether ACM alone, in the absence of serum, would be sufficient to induce phagocytosis (although serum is only added to cells for the 3-hr duration of the engulfment assay). After adding the labeled synaptosomes, we performed confocal imaging (Figure 2F) and fluorescence-assisted cell sorting (FACS) to analyze the percentage of cells with PhrodoRed fluorescence. While synaptosome phagocytosis was greatest with serum present, human astrocytes incubated with ACM alone also significantly engulfed synaptosomes (Figures 2D–2F; percentage of PhrodoRed-positive cells: no synaptosome,  $1.0\% \pm 0.4\%$ ; with synaptosomes,  $4.8\% \pm 1.1\%$ ; with synaptosomes and serum,  $13.0\% \pm 3.3\%$ ;  $n = 3$  patients).

### Human Astrocytes Have Larger Territories and More Complex Morphologies than Rat Astrocytes In Vitro

Previous studies found that human astrocytes are larger than mouse and rat astrocytes in vivo (Oberheim et al., 2006, 2009). We wondered whether the size of human astrocytes is determined by intrinsic qualities of the astrocytes or via non-cell-autonomous mechanisms. To distinguish between these possibilities, we purified human and rat astrocytes by immunopanning, cultured them in identical serum-free culture media at comparable densities, and quantified the length of processes at 5–6 days in vitro (d.i.v.). We found that the total arborization length of human astrocytes was significantly greater than that of rat astrocytes in vitro (Figure S1) (average total human astrocyte process length:  $566 \pm 56 \mu\text{m}$ ; for human astrocytes:  $n = 20$  cells from five patients; for rat astrocytes:  $n = 10$  cells from six animals). Additionally, human astrocytes had, on average, almost twice the number of branches as rat astrocytes (human:  $8.5 \pm 1.1$ ; rat:  $4.5 \pm 0.5$ ) and territories 8.6 times the size of those of rat astrocytes: for human,  $(51 \pm 13) \times 10^3 \mu\text{m}^2$ ; for rat,  $(5.9 \pm 2.9) \times 10^3 \mu\text{m}^2$ . These data are consistent with in vivo measurements (Oberheim et al., 2009). Therefore, intrinsic differences contribute, at least in part, to the larger size of human astrocytes compared with rat astrocytes.

### Human Astrocytes Exhibit Distinct Calcium Response Properties Compared with Murine Astrocytes


Mouse and rat astrocytes respond to sensory input, synaptic release of glutamate, and extracellular ATP via elevations of intracellular calcium concentrations (Ding et al., 2013; Nimmerjahn et al., 2009; Paukert et al., 2014; Schummers et al., 2008).

These calcium transients have been proposed to be important for synaptic transmission, plasticity, and regulation of blood flow (Aguilhon et al., 2010; Hamilton and Attwell, 2010; Khakh and McCarthy, 2015; Nedergaard, 1994; Parpura et al., 1994). To investigate whether human astrocytes also respond to these extracellular cues, we performed calcium-imaging studies using the calcium-sensitive dye Fluo-4 AM on acutely purified human astrocytes. We found that, at both fetal and adult stages, human astrocytes responded robustly to ATP (Figures 3A, 3C, and 3E).

Next, we wondered whether human astrocytes would respond to glutamate. Despite previous observations that adult mouse astrocytes are unresponsive to glutamate receptor agonists (Sun et al., 2013), we found that adult human astrocytes readily responded to glutamate at sub-micromolar concentrations (Figures 3A–3E). To better understand the glutamate sensitivity of human astrocytes, we sought to identify which receptor might mediate glutamate responsiveness. We found low, but non-negligible, levels of mGluR5 mRNA in human astrocytes (FPKM =  $0.34 \pm 0.06$ ,  $n = 9$ , age range = 21–63 years), which is consistent with previous reports in human and mouse (Sun et al., 2013). First, we tested the effect of the group I mGluR agonist, dihydroxyphenylglycine (DHPG), on adult human astrocytes and found that DHPG administration reliably induced robust calcium responses (Figures 3F and 3G). To further dissect which group I mGluR mediates the glutamate response in adult human astrocytes, we applied the mGluR5-specific antagonist 2-methyl-6-phenylethynyl-pyridine (MPEP) to adult human astrocytes prior to the addition of glutamate. The administration of MPEP completely blocked any glutamate response (Figures 3H and 3J). It is noteworthy that human astrocytes still responded robustly to ATP stimulation in the presence of MPEP, indicating that MPEP does not interfere with calcium storage or release machinery in astrocytes (Figure 3J). After washout of MPEP for 30 min, the cells again robustly responded to glutamate (Figures 3I and 3K).

To assess the development of glutamate responsiveness in human astrocytes, we compared the glutamate responses of fetal human and rat astrocytes. Astrogenesis commences shortly before weeks 17–20 of gestation in humans and embryonic day (E) 18.5 in rats. Therefore, we considered these stages developmentally comparable in these two species. In contrast to rat fetal (E18.5) astrocytes, which readily responded to glutamate at sub-micromolar concentrations, fetal human astrocytes did not respond to glutamate at micromolar or even millimolar concentrations (percentage of cells responding to 30 nM, 300 nM, 800 nM, 1.7  $\mu\text{M}$ , and 3  $\mu\text{M}$  glutamate: 5%, 70%, 70%, 100%, and 100%, respectively, for rat fetal astrocytes; 0% at all concentrations tested for human fetal astrocytes;  $n = 20$  cells per condition; Figure 3L). Therefore, human astrocytes likely acquire the ability to respond to glutamate at a stage later than mid-gestation.

Although adult human astrocytes responded to both ATP and glutamate, closer examination of the calcium transients indicated that ATP and glutamate elicited responses with distinct temporal patterns. Independent of concentration, glutamate stimulation consistently produced a synchronous “all or none” rise in intracellular calcium levels across most astrocytes that then quickly decreased (within 10 s) to baseline levels (Figures



### Figure 3. Calcium Responses of Human Astrocytes In Vitro

(A) Representative images of calcium responses to 3  $\mu\text{M}$  ATP (top) and glutamate (bottom) of mature human astrocytes (21 years old; yo). Time is labeled in each image. Arrows point to two cells showing calcium fluctuations in response to ATP.

(B and C)  $\Delta F/F$  of all cells in (A). Each  $\Delta F/F$  trace represents a single cell.

(D) Average number of calcium fluctuations in human astrocytes after exposure to ATP or glutamate.  $n = 11$  movies. Concentrations of ATP and glutamate: 30 nM–3  $\mu\text{M}$ . \* $p < 0.05$ , two-tailed Wilcoxon rank-sum test. Error bars represent mean  $\pm$  SEM.

(E) Percentage of cells responding to ATP or glutamate at different concentrations. Cells with  $\Delta F/F > 5\%$  are counted as responsive.  $n = 3$ –5 movies per condition. Each movie has 5–20 cells. Data were collected from two patients 21 and 26 years of age, respectively. Error bars represent mean  $\pm$  SEM.

(F) Fluo-4 AM fluorescence of adult human astrocytes before and after addition of DHPG (200  $\mu\text{M}$ ).

(G)  $\Delta F/F$  of adult human astrocytes before and after addition of DHPG.

(H) Fluo-4 AM fluorescence of adult human astrocytes before and after addition of 3  $\mu\text{M}$  glutamate in the presence of 25  $\mu\text{M}$  MPEP.

(I) Fluo-4 AM fluorescence of adult human astrocytes before and after addition of 3  $\mu\text{M}$  glutamate after washing out MPEP for 30 min.

(J)  $\Delta F/F$  of adult human astrocytes before and after addition of 3  $\mu\text{M}$  glutamate and 3  $\mu\text{M}$  ATP in the presence of 25  $\mu\text{M}$  MPEP. Arrowheads represent addition of glutamate or ATP.

(K)  $\Delta F/F$  of adult human astrocytes before and after addition of 3  $\mu\text{M}$  glutamate after washing out MPEP.

(L) Representative images of calcium responses to 3  $\mu\text{M}$  glutamate in fetal human (17–20 gw, left) and rat (E18.5, right) astrocytes. Scale bar, 50  $\mu\text{m}$ .



3B and 3C). In contrast, the onset of calcium response to ATP stimulation was asynchronous among the astrocytes in the imaging chamber. After the initial calcium elevation, some cells exhibited fluctuations in calcium concentrations that lasted for minutes after initial ATP stimulation (Figures 3A and 3C). It remains unclear whether ATP and glutamate elicit calcium responses with similar temporal dynamics in murine astrocytes.

### Transcriptomes of Acutely Purified Human Neurons, Astrocytes, Oligodendrocytes, Microglia, and Endothelial Cells

The human brain is composed of neurons, glia (astrocytes, oligodendrocytes, microglia), and vascular cells. Each of these cell types has a distinct role to play in the nervous system, and each can be characterized by a unique molecular repertoire necessary for cell-specific functions. To construct a transcriptome database of purified human neurons, astrocytes, oligodendrocytes, microglia, and endothelial cells, we acutely isolated each cell type by immunopanning (Figure 1), extracted RNA, and performed RNA-seq (31 million  $\pm$  4 million pair-end 150-bp reads per sample). RNA was collected immediately at the conclusion of the purification protocol to best reflect the true *in vivo* molecular profiles. Although we took great effort to minimize the duration of time between tissue harvest and RNA collection, gene expression may change during dissociation and purification. Therefore, we collected RNA from cortical tissue immediately after resection and compared these data to RNA collected from the same cortical sample after dissociating the tissue and letting it sit for the duration of a typical purification timeline (within about 5 hr from tissue resection). We found a strong correlation between the two samples, with only 1.49% of the genes changing more than 4-fold in either direction during the tissue dissociation protocol (294/19,713 genes). Those genes on this small list largely constitute immediate early genes from the acute macrophage response to injury (Tables S2 and S5). We cross-referenced this list with our downstream analyses to ensure that these genes did not confound the interpretation of our data.


To first assess the purity of our immunopanning-isolated cell samples, we probed the transcriptome data for expression of well-known cell-specific genes for astrocytes (e.g., *GFAP*, *ALDH1L1*, *SOX9*, and *AQP4*), neurons (e.g., *VGLUT1*, *STMN2*, *SYT1*, *SYN1*), oligodendrocytes (e.g., *PLP1*, *MOG*, *SOX10*, and *MBP*), microglia/macrophages (e.g., *C1QA*, *CX3CR1*, *CCL3*, and *TNF*), and endothelial cells (e.g., *CLDN5*, *ELTD1*, *ITM2A*, and *ESAM*) (Figures 4A and 4B; Table S4). The expression of these classical cell-specific markers each demonstrated definitive cell-type selectivity in their corresponding cell types with undetectable or low levels of expression by other cell populations not known to express these markers (Figures 4A and 4B; Table S3) ( $N_s = 12, 1, 5, 3,$  and  $2$  patients for astrocytes, neurons, oligodendrocytes, microglia, and endothelial cells, respectively). To assess the extent of individual differences between human patients, we examined the correlation between replicates of identical cell types across different patients ( $n = 23$ , age range = 8–63 years) and found high correlations ( $0.83 \pm 0.02$ , Spearman correlation) between cell-type replicates and low correlations ( $0.34 \pm 0.013$ ) among samples of differing

cell types (Figure 4C). Next, we specifically examined the variation between replicates of human astrocyte samples ( $0.89 \pm 0.05$ ). To determine the degree to which variation among these samples was due to variability in the panning procedure, as opposed to inherent inter-patient differences, we performed RNA-seq on four whole cortex samples (specimens that had not been immunopanned) from separate individuals. We found almost identical correlations between whole-cortex samples ( $0.89 \pm 0.08$ ) and what we observed in the immunopanned astrocyte samples, suggesting that the variance across astrocyte samples can largely be attributed to small, inherent levels of variation across individuals. As expected, mouse astrocyte samples from an inbred strain (C57BL6) demonstrated a slightly higher correlation ( $0.94 \pm 0.04$ ) than samples from different human patients.

Additionally, we performed unsupervised hierarchical clustering and found that samples of identical cell types clustered together rather than clustering by patient (Figure S4). Together, these data suggest that transcriptome differences between cell types are not obscured by individual genetic and/or environmental variations between subjects. To disseminate these human cell-type-specific gene expression data to the greater neuroscience community, we generated a user-friendly searchable online database of the gene expression profiles of all acutely purified human cell types (<http://www.brainmaseq.org>).

### Transcriptome Differences between Human APCs and Mature Astrocytes

The molecular and cellular changes that occur during astrocyte maturation are poorly understood. To examine gene expression changes associated with astrocyte maturation, we compared transcriptomes of APCs and mature human astrocytes. Whereas a large number of genes were expressed both by APCs and mature astrocytes (Figures 5A and 5B; fetal,  $n = 6$  patients; mature,  $n = 12$  patients), a substantial contingent ( $>2,000$ ) of genes were differentially expressed (DE) between the two stages (Table 1). For example, APCs expressed high levels of proliferative genes such as *MKI67* and *TOP2A*, confirming their immature properties (Figures 5B and 5F). In contrast, a number of genes were only expressed by mature astrocytes, such as gap junction genes *GJA1* and *GJB6*, the human-neuropsychiatric-disorder-associated gene *AGXT2L1*, and a Wnt pathway inhibitory factor *WIF1*. These genes provide new markers that could be used for the discrimination of developing and mature human astrocytes. To take an unbiased approach to find pathways unique to APCs and mature astrocytes, we identified all significantly DE genes between the two developmental stages (using volcano plots with cutoffs at  $p = 0.01$  and 4-fold ratios; Figure S5C) and used these DE genes to run a Gene Ontology (GO) term analysis with DAVID bioinformatics resources (Huang et al., 2009). As astrocytes matured, we found a significant downregulation of cell-cycle genes and a significant upregulation in genes involved with nerve impulse transmission, cell-cell signaling, fatty acid metabolism, cell adhesion, and ion homeostasis (a full list of GO terms and DE genes is given in Tables S5 and S6). We performed identical volcano plot and DAVID analyses for subsequent comparisons between cell types and species (Figures S5B–S5E; Tables S5 and S6).



**Figure 4. RNA-Seq Transcriptome Profiling of Acutely Purified Human Neurons, Astrocytes, Oligodendrocytes, Microglia/Macrophages, and Endothelial Cells**

(A) Expression of classic cell-type-specific genes by acutely purified mature human neurons, astrocytes, oligodendrocytes, microglia/macrophages, and endothelial cells. Data represent SD from mean across rows.  $N_s = 12, 1, 5, 3,$  and  $2$  patients for astrocytes, neurons, oligodendrocytes, microglia, and endothelial cells, respectively. Age range = 8–63 years.

(B) Representative examples of the expression of cell-type-specific genes by acutely purified cell samples. Error bars represent mean  $\pm$  SEM.

(C) Spearman correlation between all fetal and mature samples. Fetal,  $n = 6$  patients. Age range = 18.0–18.5 gw. Mature,  $n_s = 12, 1, 5, 3,$  and  $2$  patients for astrocytes, neurons, oligodendrocytes, microglia, and endothelial cells, respectively. Age range = 8–63 years.

(D) Scatterplot of human and mouse astrocyte gene expression.  $\rho$  represents the Spearman correlation coefficient. Venn diagram indicates the number of astrocyte-enriched genes (>4-fold, FPKM > 5) for human and mouse astrocytes. Average expression data from all samples of mature human astrocytes were used for this analysis. Human,  $n = 12$  patients. Mouse,  $n = 4$  batches of astrocytes, each comprising three mouse brains combined. Age range of human patients = 8–63 years. Age range of mice = 1–9 months. Fold enrichment in astrocytes is calculated as FPKM in astrocytes divided by the average FPKM in all the other cell types (neurons, oligodendrocytes, microglia, and endothelial cells).

(E) Differences in gene expression between purified human and mouse astrocytes measured using qPCR. Fold changes reflect enrichment of specific genes in human tissues. Error bars represent mean  $\pm$  SEM.

(F) In situ hybridization performed on temporal lobe cortices from healthy patients (separate source from the RNA-seq samples). Probes were designed against LRRC3B and GPR98 (cyan), and sections were counterstained against the astrocyte-specific transcription factor, SOX9 (magenta). Scale bars,  $100 \mu\text{m}$  zoomed out; scale bars in insets,  $50 \mu\text{m}$ .



**Table 1. Top Gene Expression Differences between APC and Mature Astrocytes**

Gene	Expression in Fetal Astrocytes	Expression in Mature Astrocytes	Fold Change
<b>Top Mature Astrocyte Genes</b>			
<i>AGXT2L1</i>	0.1	238.6	1,988.6
<i>S100A1</i>	0.1	84.9	849.4
<i>SLC14A1</i>	0.1	29.3	334.6
<i>TMEM176A</i>	0.1	27.1	270.6
<i>TMX2</i>	0.1	25.9	259.1
<i>HHATL</i>	0.1	24.9	237.6
<i>PADI2</i>	0.1	26.8	225.3
<i>TLR4</i>	0.1	18.7	212.9
<i>HSD17B6</i>	0.1	20.5	204.6
<i>CHI3L1</i>	0.1	15.2	172.2
<i>NUDT3</i>	0.1	13.5	135.2
<i>FBXO2</i>	0.1	10.8	115.1
<i>NTSR2</i>	0.1	10.8	107.5
<i>ALDH1L1</i>	0.1	10.8	82.5
<i>ALDOC</i>	4.8	371.0	77.6
<i>SLC1A2</i>	34.5	1,987.9	57.6
<i>RYR3</i>	0.3	16.0	55.0
<i>GABRA2</i>	0.9	46.1	51.2
<i>CPE</i>	16.0	657.2	41.0
<i>GLUL</i>	21.0	826.9	39.4
<b>Top APC Genes</b>			
<i>HIST1H2AI</i>	57.9	0.1	578.9
<i>HIST1H3E</i>	51.0	0.1	509.6
<i>HIST1H3B</i>	176.6	0.6	292.1
<i>HIST1H1B</i>	23.4	0.1	234.0
<i>PPDPF</i>	26.4	0.2	119.3
<i>TPX2</i>	22.0	0.2	93.9
<i>NUSAP1</i>	27.4	0.3	90.4
<i>HIST1H2AC</i>	39.0	0.5	76.8
<i>HIST2H2AC</i>	68.9	1.3	53.2
<i>TNC</i>	26.6	0.6	45.5
<i>KIF15</i>	17.2	0.5	32.7
<i>HIST1H1A</i>	45.9	1.5	30.7
<i>HIST1H2BC</i>	215.2	13.2	16.3
<i>FABP5</i>	23.9	1.5	15.5
<i>DTYMK</i>	10.3	0.7	14.8
<i>RAB11B</i>	10.9	0.8	13.5
<i>HES6</i>	21.6	2.1	10.3
<i>LRIG3</i>	13.3	1.4	9.6
<i>E2F5</i>	10.1	1.4	7.2
<i>MPPED2</i>	16.0	2.3	7.0

Genes are ranked by fold change in expression between APC and mature stages. The top and bottom 20 genes are listed. Genes with FPKM <10 in both APC and mature stages are excluded to highlight highly expressed genes that are more likely to be useful markers for APC or mature astrocytes. All genes have >2-fold enrichment in astrocytes over other cell types.

RNA-seq data from mouse astrocytes were previously collected by FACS from transgenic lines (Zhang et al., 2014), we collected new mouse astrocyte samples using an identical procedure to our human astrocytes, including the use of the monoclonal HepaCAM antibody for immunopanning. To ensure that HepaCAM-panned mouse astrocytes were comparable to mouse astrocytes purified via our previous method of FACS sorting ALDH1L1-EGFP mice, we collected and compared age-matched FACS-sorted ALDH1L1-EGFP astrocytes with HepaCAM-panned astrocytes. Again, using volcano plots with DE cutoffs of  $p = 0.01$  and 4-fold ratio changes, we could detect only 55 total DE genes (0.3%) between the two different methods of mouse astrocyte purification (Figure S5A; Table S6). Then, we compared the RNA-seq-based transcriptomes of human and mouse HepaCAM-purified astrocytes and found significant overlap including all of the classic astrocyte-specific genes (e.g., *GFAP*, *ALDH1L1*, *GLUL*, *AQP4*, *SLC1A2*, and *SLC1A3*). Using a threshold of 4-fold enrichment (calculated as expression in astrocytes divided by the average expression level in all other cell types—neurons, oligodendrocytes, microglia, and endothelial cells), 52% of mouse astrocyte-enriched genes were similarly enriched in human astrocytes. Conversely, only 30% of human astrocyte-enriched genes were identified as astrocyte enriched in mice (Figure 4D; Table 2) (human,  $n = 12$  patients; mouse,  $n = 4$  batches of astrocytes each made from three mouse brains combined). These trends were consistent across a range of thresholds from 2-fold to 50-fold (data not shown). Next, we examined whether human-mouse differences in gene expression were more prominent in specific cell types. We found almost identical transcriptome-level differences between human and mouse in each of the brain cell types that we examined, and these values were also comparable to those for non-neural tissues such as liver and muscle cells (data not shown).

One further concern about interpreting interspecies differences is that the human data are exclusively derived from temporal lobe, in comparison to whole-cortex data from mouse. To address this potential confounder, we utilized the Allen Brain Atlas dataset (Hawrylycz et al., 2012) to determine all genes that are enriched in temporal lobe versus other cortical areas and found no overlap with our candidate human versus mouse enriched genes (Table 2; Table S7). Despite the homology between human and mouse genomes, there are a number of human genes and long non-coding RNAs without mouse orthologs. We found that some of these human-specific genomic elements are expressed in cell-type-specific patterns (Table S8). Further investigation of these candidates may provide novel insights into new properties that have evolved from mouse to human in various CNS cell types.

#### Validation of DE Genes from Human and Mouse


Next, we examined the validity of our RNA-seq transcriptome datasets by orthologous methods. First, we examined a subset of genes identified by our RNA-seq data to be DE genes from human and mouse cells using PCR on cDNA generated from whole mouse and human brain samples. PCR amplification of all of the genes we examined (*WIF1*, *MRV11*, *APOC2*, *GLDN*, *GPR98*, *PMP2*, *CNDP1*, and *MAG*) showed species-selective



**Table 2. Top Astrocyte-Enriched Genes Shared by Humans and Mice and Species-Specific Astrocyte Genes**

Human and Mouse Astrocyte-Specific Markers			Genes in Human Astrocytes Not Found in Mouse Astrocytes				Genes in Mouse Astrocytes Not Found in Human Astrocytes			
Gene Symbol	Human Expression	Mouse Expression	Gene Symbol	Human Astrocyte Expression	Mouse Astrocyte Expression	Fold Enrichment in Human Astrocytes/ Mouse Astrocytes	Gene Symbol	Human Astrocyte Expression	Mouse Astrocyte Expression	Fold Enrichment in Mouse Astrocytes/ Human Astrocytes
GJA1	560.3	351.5	WIF1*	66.9	0.1	668.7	Pla2g7*	0.2	138.0	866.5
SLC39A12	79.3	88.6	PMP2	80.1	0.1	663.8	H3f3a	0.1	56.2	561.9
GJB6	106.0	88.8	GSTM2*	43.6	0.1	331.1	Thrsp	0.1	36.1	360.6
BMPR1B	27.0	24.7	LRRC3B*	29.7	0.1	245.0	Itih3*	0.1	28.5	285.2
SLC14A1	29.3	37.8	HSD17B6*	20.5	0.1	204.6	Lxn*	0.2	61.2	265.0
PAMR1	23.1	23.8	FAM198B*	20.0	0.1	198.9	Vcam1*	0.2	45.6	226.6
RGS20	17.9	52.6	PNMA1	26.1	0.1	187.7	Nnat	0.1	21.8	218.3
FGFR3	38.4	182.4	MRVI1*	18.3	0.1	183.2	Tnfrsf19	0.2	37.9	217.6
MGST1	72.4	18.1	RYR3*	16.0	0.1	160.0	Ngef	0.1	22.1	211.9
SLC4A4	97.6	199.4	GSTM3	19.3	0.1	144.1	Snpc5	0.1	19.6	195.7
AQP4	450.0	198.4	APOC2	15.3	0.1	137.4	Chst2	0.3	57.3	195.0
F3	126.1	112.6	RGN	18.1	0.1	133.5	Magoh	0.1	18.0	180.0
MLC1	15.3	242.3	ACBD7*	21.6	0.2	132.3	Romo1	0.1	16.9	169.3
ELOVL2	41.2	34.2	RGR*	12.5	0.1	115.6	Akr1b10	0.1	16.2	161.9
SLC1A2	1,987.9	1,087.4	AMY2B	11.5	0.1	114.5	Fam176a	0.1	16.4	160.5
CLU	384.2	769.3	NDUSF5	27.3	0.2	112.3	Il18*	0.7	107.8	154.8
ACSBG1	19.6	140.8	STOX1*	18.1	0.2	107.3	Pou3f3	0.1	15.1	150.6
TTYH1	16.4	360.5	GPR98*	50.3	0.5	97.1	Fam173a	0.1	16.2	145.9
ATP1B2	468.7	550.7	AADAT*	10.7	0.1	79.2	Rnf5	0.2	24.6	142.9
SOX9	50.5	57.7	CALCOCO2	10.5	0.1	75.4	tspan2	0.1	13.5	134.4

Shared astrocyte genes are expressed in both human and mouse astrocytes (FPKM > 0.5) and enriched by at least 4-fold over the average expression in neurons, oligodendrocytes, microglia/macrophages, and endothelial cells. Species-specific astrocyte genes have FPKM values <1 in astrocytes from the non-enriched species, and those that are also astrocyte enriched are indicated with asterisks. Top 20 genes are listed for all categories (ranked by fold enrichment).



### Figure 6. Acutely Purified Human Astrocytes Display Resting Gene Profiles

(A) The Zamanian et al. (2012) dataset was used to identify the top 30 reactive astrocyte genes upregulated following middle cerebral artery occlusion (MCAO), LPS infection, or both. Expression of this suite of “reactive” genes was then probed in various astrocyte samples. The white dots in the Violin plots represent the median. Thick vertical black lines represent interquartile ranges. Horizontal dashed lines indicate the median in normal human astrocytes for each condition. Ns = 12, 4, 3, and 3 for normal, epilepsy, tumor, and serum-expanded astrocytes, respectively. (B) Representative images of acutely purified human astrocytes grown in culture for 7 days in the absence (left) or presence (right) of serum. Scale bars, 20  $\mu$ m.

### Reactive Changes of Human Astrocytes in Neurological Disorders

Previous methods of enriching human astrocytes (the McCarthy de Vellis, or MD method) in serum-based cultures involve long-term exposure to a non-physiological environment, and serum exposure in culture has been shown to induce irreversible reactive changes in astrocytes (Foo et al., 2011; Zamanian et al., 2012). Furthermore, astrocytes grown in vitro via serum-selection methods demonstrate unusual polygonal fibroblast-like morphologies that are not present in vivo, whereas our acutely purified astrocytes grown in serum-free medium exhibit extensive process-bearing morphologies (Figures 1 and S3). To test whether exposure to serum could elicit morphological changes in acutely purified human astrocytes, we added serum to our cultures of immunopanned human astrocytes and quickly noted a transition from a process-bearing morphology to the polygonal fibroblast-like morphology typically found in serum-cultured cells (Figure 6B).

expression patterns as predicted by RNA-seq (Figure S6). Next, we used qPCR to validate a subset of the top candidates for DE genes between fetal versus adult human and mouse versus human astrocytes and found consistent agreement with our RNA-seq findings in each gene that we investigated (Figures 4E and 5G).

Additionally, we used in situ hybridization to examine the spatial expression of two human astrocyte-enriched genes on human brain sections. We combined fluorescence in situ hybridization with probes designed against two of the top human-specific astrocyte candidates, *GPR98* and *LRR3B*, and immunofluorescence with an antibody against SOX9, an astrocyte-specific nuclear protein. We found strong colocalization between our RNA probes and SOX9 staining, providing evidence that *GPR98* and *LRR3B* are both highly enriched in human astrocytes (Figure 4F), as predicted by the RNA-seq data.

To examine whether the transcriptome profiles of acutely purified human astrocytes more closely resembled resting or reactive astrocytes, we probed the RNA-seq dataset of human astrocytes for reactive astrocyte genes. Previous reports of mouse reactive astrocytes have described two distinct types of reactive phenotypes: a bacterial lipopolysaccharide (LPS)-infection-induced “A1” phenotype and an ischemia-induced “A2” phenotype (Zamanian et al., 2012). We cross-referenced the top 30 genes that have been shown in astrocytes to be induced by ischemia alone, LPS alone, or both injury models with our acutely purified human astrocytes. We found only low expression of these reactive astrocyte markers in our preparations of acutely purified human astrocytes (Figure 6A, n = 12 patients). In contrast, we found that human astrocytes obtained

from the Zamanian et al. (2012) dataset were highly expressed in our preparations of acutely purified human astrocytes (Figure 6A, n = 12 patients). In contrast, we found that human astrocytes obtained

using serum selection methods from two commercial sources, Lonza and ScienCell (Lonza astrocytes,  $n = 1$ ; and ScienCell astrocytes,  $n = 2$ ) (Malik et al., 2014), expressed high levels of these reactive astrocyte markers.

What are the transcriptome-wide changes that affect human astrocytes in neurological diseases? Numerous postmortem immunofluorescence studies of patients with a variety of neurological conditions have demonstrated upregulation of a single reactive astrocyte marker, GFAP (Beach and McGeer, 1988), but a transcriptome level examination of reactive changes in human astrocytes has not yet been performed. To examine disease-induced gene expression changes in human astrocytes, we acutely purified astrocytes from brain samples taken from regions involved in epilepsy and glioblastoma. Then, we performed RNA-seq to compare transcriptomes of disease-affected astrocytes with healthy human astrocytes. Much like in serum-cultured astrocytes, we found significant increases in the expression of reactive astrocyte genes in disease-affected astrocytes (Figure 6A; Table S6). These results support numerous morphological observations that human astrocytes undergo robust reactive changes in epilepsy and glioblastoma.

## DISCUSSION

We have developed a method to acutely purify human astrocytes and to grow these cells in serum-free conditions *in vitro*. This method allowed us, for the first time, to directly investigate the functions of human astrocytes. Like mouse astrocytes, we found that human astrocytes strongly promote neuronal survival, synapse formation, and engulf synapses. We also found that human astrocytes retain their larger size *in vitro* and that adult human astrocytes demonstrate robust mGluR5-mediated calcium responses to glutamate stimulation. Next, we performed RNA-seq transcriptome profiling of purified human neurons, astrocytes, oligodendrocytes, microglia, and endothelial cells and established a database that serves as a road map for understanding cell-type-specific gene function in the human brain. By comparing human and mouse astrocyte transcriptome profiles, we found large numbers of genes shared by astrocytes of both species, as well as a gene expression signature specific to human astrocytes. We further demonstrated that acutely purified human astrocytes from normal brain samples exhibit resting astrocyte gene expression profiles and that epilepsy and glioblastoma robustly induce reactive changes in human astrocytes.

### Application of New Immunopanning Purification Protocol for Studies of Human Brain Cell Types

The ability to purify and culture human astrocytes opens many new avenues of investigation. The vast majority of drug candidates for neurological disorders succeed in mouse model studies but fail in human clinical trials (Kola and Landis, 2004). Thus, there is a strong need for human cell culture models for disease mechanism studies and drug testing. Acutely purified human astrocytes can be cultured for several weeks *in vitro* and maintain functional properties at these later time points (Figures S1 and S3). It should be noted, however, that the transcriptome profiles of “resting” human astrocytes are derived from acutely purified astrocytes and not those in culture.

Since astrocytes have been found to be key players in many neurological disorders, improvement of human astrocyte culture methods may lead to new discoveries in the treatment of neurological disorders. Previously developed serum-based culture methods generate human astrocytes that express reactive astrocyte genes and are already in a disease-like state. Therefore, it is difficult to identify additional disease-induced changes above the background reactive changes with these cells. The data presented here of acutely purified human astrocytes provide a baseline by which to compare changes in gene expression in diseased versus healthy brains.

### Calcium Responses in Human Astrocytes

The glutamate response of adult human astrocytes is surprising, in light of a recent study on mouse astrocytes that reported that only young mouse astrocytes respond to a glutamate receptor agonist (Sun et al., 2013). Like Sun et al., we found low mRNA levels of mGluR5 in adult human astrocytes, but we pursued pharmacological studies to examine whether this low expression might mediate the robust glutamate response of adult human astrocytes. Our pharmacological experiments suggest that human astrocyte glutamate signaling occurs via mGluR5. The difference in the ability of adult human and mouse astrocytes to respond to extracellular glutamate suggests that adult human astrocytes may have evolved an improved capability to detect and potentially respond to synaptic activity. Whether such a difference contributes to the unique cognitive abilities of humans remains an intriguing question.

### Two Distinct Stages of Human Astrocyte Development

The assembly of neural circuits and the maturation of the brain involve precisely orchestrated changes of neurons and glia. The distinct characteristics of neural progenitors and mature neurons, OPCs, and mature oligodendrocytes have been well defined. Astrocyte maturation, however, is much less understood. Mouse astrocytes are generated from radial glial cells during late embryogenesis and early postnatal development, and they continue to divide and populate the brain during the first 2 postnatal weeks (Ge et al., 2012; Tien et al., 2012). After proceeding through a poorly understood maturation process, astrocytes lose their abilities to divide, develop elaborate processes that ensheath synapses, and surround blood vessels to form the blood-brain barrier. Investigating how molecular profiles of human astrocytes mature throughout development is critical not only for understanding brain development but also for gaining insights into pathologies where astrocytes acquire fetal phenotypes (i.e., glioblastoma) (Figure 5).

In this study, we found that human astrocytes exist in at least two distinct states: a fetal highly proliferative APC state and a postnatal (childhood and adult), non-proliferative, mature state, bearing much similarity with OPCs and mature oligodendrocyte stages. As has been previously observed in mouse astrocytes (Ge et al., 2012; Tien et al., 2012), astrocytes isolated from fetal human brains had a vastly higher proliferative ability than mature astrocytes. We did not observe proliferation (or the presence of mitotic genes) in astrocytes isolated from older human brains, even in the presence of high levels of the mitogen HBEGF. Our transcriptome profiling of fetal, juvenile, and adult

human astrocytes provides a molecular description of the numerous changes that occur during astrocyte maturation. As human astrocytes mature, they turn on genes encoding synapse-inducing proteins (e.g., SPARCL1), gap junction proteins (e.g., GJA1 and GJB6), proteins for neural transmitter recycling (e.g., SLC1A2, SLC1A3, and GLUL), secreted proteins (e.g., CLU and SPON1), a thyroid hormone metabolism enzyme (DIO2), and a variety of transmembrane receptors (e.g., FGFR3 and GABRA2). We cross-referenced a published dataset of human brain gene expression across developmental stages (Kang et al., 2011) and found that mature astrocytes are hardly present prior to birth in most of the regions in the human brain. The vast majority of mature astrocyte markers rapidly increase in expression soon after birth, peaking around 6 months to 1 year of age. Interestingly, changes of synapse density in the developing human cortex follow an identical time course (Huttenlocher, 1979; Huttenlocher et al., 1982). Furthermore, a phagocytic receptor required for astrocyte-mediated synapse elimination, MERTK (Chung et al., 2013), is not expressed by fetal astrocytes but later turns on and reaches peak levels around 6 months to 1 year of age. Concurrently, synapse density begins to decline after 1 year of age in humans (Huttenlocher, 1979; Huttenlocher et al., 1982). The precise temporal correspondence of human astrocyte maturation and synapse density, as well as previously reported roles of astrocytes in synapse formation and elimination, suggests that astrocytes may trigger maturation of neural circuitry. Consistent with this notion, a number of genes associated with neurodevelopmental disorders such as autism and schizophrenia are enriched in astrocytes (data not shown). An important goal for future study is to understand the intrinsic or extrinsic signaling mechanisms that trigger astrocyte maturation. Alterations in the timing of this maturation could potentially underlie neurodevelopmental disorders.

### Astrocyte Transcriptome Changes in Epilepsy and Glioblastoma

Our immunopanning technique allowed us to purify and profile astrocytes from diseased environments. At the transcriptomic level, our data corroborate numerous morphological observational studies of astrocytes in these conditions. We found over 400 DE genes between healthy astrocytes and those from sclerotic hippocampus, and similar numbers from those in glioblastoma. A careful examination of the gene expression changes in these disease states may potentially generate new insights into the role of astrocytes in these diseases.

### Species Differences in Astrocyte Properties

To compare the transcriptomic landscape of human and mouse astrocytes, we began by performing an unbiased genome-wide comparison of the two species. We found over 600 human astrocyte-enriched genes that were not similarly enriched in mouse astrocytes. Some of these genes may provide insight into the special physiological and functional properties of human astrocytes. For example, we found that the ryanodine receptor 3 (Ryr3) is highly expressed by human astrocytes but not by mouse astrocytes. Ryr3 is a calcium-permeable ion channel located on the membrane of the endoplasmic reticulum (ER),

where calcium is stored inside cells. It is activated by elevation of the cytoplasmic calcium concentration and releases calcium from the internal store in a process called “calcium-activated calcium release” in muscle cells (Gordienko and Bolton, 2002). The presence of Ryr3 in human astrocytes might amplify the elevation of intracellular calcium concentration and allow fast propagation of calcium transients along astrocyte processes by releasing calcium from spatially discrete ER compartments. We also found human-specific expression of MRV11, a protein that binds the IP3 receptor and regulates intracellular calcium stores (Figure S6). These findings are consistent with a study reporting differences in the propagation of calcium transients between human and mouse astrocyte networks (Oberheim et al., 2009).

As the evidence accumulates for the importance of glia—and, particularly, astrocytes—in health and disease, the purification method and transcriptome datasets presented here will be a valuable resource for investigating the biology of human astrocytes and for investigating new treatment approaches for neurological disorders.

## EXPERIMENTAL PROCEDURES

### Human Tissue

Human brain tissue was obtained with informed consent under a protocol approved by the Stanford University Institutional Review Board. Juvenile and adult human brain tissue samples were obtained from surgeries for treating epilepsy and tumors. All experiments described in this study, except the reactive astrocyte gene induction studies (Figures 5 and 6, epilepsy, tumor, and tumor-peripheral data), were performed with, in general, normal temporal lobe cortex resected in order to access deeper hippocampus areas involved in epilepsy. The cortical tissue was determined as normal by EEG and MRI. For assessment of reactive astrocyte gene expression, we used sclerotic hippocampus specimens involved in epileptic foci and cortical specimens from glioblastoma core and peripheral regions, defined as contrast enhancing and non-contrast-enhancing regions, respectively. Tissue was immersed in 4°C Neurobasal medium (Thermo Fisher Scientific) and transferred to the lab for tissue dissociation within 1 hr after resection. Fetal human brain tissue was obtained with informed consent following elective pregnancy termination under a protocol approved by the Stanford University Institutional Review Board. Tissue was immersed in 4°C RPMI medium and transferred to the lab for tissue dissociation within 5 hr after the procedure.

### Vertebrate Animals

All procedures involving animals were conducted in conformity with Stanford University guidelines that are in compliance with national and state laws and policies. We used mice for RNA-seq transcriptome profiling, in situ hybridization, and PCR validation experiments of the RNA-seq data. We used rats for in vitro experiments, including morphology measurement and calcium imaging, because of higher yield and consistency of rat astrocyte culture compared to mouse astrocyte culture.

### Purification of Mature Human Astrocytes, Neurons, Oligodendrocytes, Microglia/Macrophages, and Endothelial Cells

For a detailed purification protocol, see the Supplemental Information. Briefly, we dissected out gray matter from mature human brain specimens, incubated the tissue in 20 U/ml papain at 34°C for 100 min, and washed the tissue with a protease inhibitor stock solution. Then, we gently triturated the tissue and then added the single-cell suspension to a series of plastic Petri dishes pre-coated with cell-type-specific antibodies and incubated for 10–30 min each at room temperature. Unbound cells were transferred to the subsequent Petri dish, while the dish with bound cells was rinsed eight times with PBS to wash away loosely bound contaminating cell types. The antibodies used include



anti-CD45 to capture microglia/macrophages, anti-GalC hybridoma supernatant to harvest oligodendrocytes, anti-O4 hybridoma to harvest OPCs, anti-Thy1 (CD90) to harvest neurons, anti-HepaCAM to harvest astrocytes, and BSL-1 to harvest endothelial cells. For RNA-seq, cell samples were scraped off the panning dish directly with Qiazol reagent (QIAGEN). For cell culture and *in vitro* experiments, astrocytes bound to the anti-HepaCAM antibody-coated dishes were incubated in a trypsin solution at 37°C for 5–10 min and gently squirted off the plate. Then, we spun down the astrocytes and plated them on poly-D-lysine-coated plastic coverslips in a Neurobasal-DMEM-based serum-free medium (the composition is detailed in [Supplemental Information](#)). We replaced half of the volume with fresh medium every 3–4 days to maintain the cultures. Fetal human astrocytes and neurons were purified in a protocol similar to the aforementioned procedure, with the modifications detailed in the [Supplemental Information](#).

#### RNA-Seq Library Construction and Sequencing

Total RNA was extracted using the miRNeasy kit (QIAGEN) under the protocols of the manufacturer. The quality was assessed by Bioanalyzer. Samples with RNA integrity number higher than 8 were used for library construction. We used the Ovation RNA-seq system V2 (Nugen 7102) to perform first and second-strand cDNA synthesis. We then used the Next Ultra RNA-seq library prep kit for Illumina (NEB E7530) and NEBNext® multiplex oligos for Illumina (NEB E7335 E7500) to perform end repair, adaptor ligation, and 5–6 cycles of PCR enrichment according to manufacturer's instructions. The quality of the libraries was assessed by bioanalyzer and qPCR and high quality libraries were sequenced by the Illumina NextSeq sequencer to obtain 150-bp paired reads. See detailed protocol in the [Supplemental Information](#).

#### RNA-Seq Read Mapping, Transcript Assembly, and Expression Level Estimation

We analyzed RNA-seq reads with the Galaxy web-platform (<http://usegalaxy.org>). The FASTQ files were first groomed using the FASTQ groomer and then mapped using TopHat2, which invokes Bowtie as an internal read mapper. The paired-end option was selected and human genome version 19 (hg19) and mouse genome version 9 (mm 9) was used as the reference genome for the human and mouse RNA-seq data, respectively. We then ran Cufflinks to assemble transcripts and estimate expression level as FPKM.

#### Calcium Imaging

Cells were incubated for 15 min with 2  $\mu$ M Fluo-4 AM (Invitrogen, F-14201) and then washed with PBS and replaced with growth medium. Images were taken at 0.7-s intervals and analyzed with ImageJ.

#### Immunocytochemistry

Cultured cells were fixed with 4% paraformaldehyde for 10 min at room temperature, permeabilized, and blocked with 10% goat serum with 0.2% Triton X-100. See the [Supplemental Information](#) for antibody details.

#### Electrophysiology

Whole-cell patch-clamp recordings from cultured RGC neurons were performed at room temperature in an isotonic saline solution. mEPSCs were recorded in tetrodotoxin (TTX) (1  $\mu$ M, Alomone Labs) from a holding potential of  $-70$  mV. Series resistance was monitored throughout the recording and was  $<20$  M $\Omega$ . Data were sampled at 50 kHz and filtered at 1 kHz. See detailed protocol in the [Supplemental Information](#).

#### Synapse Formation Assay

We purified rat RGCs by sequential immunopanning to greater than 99% purity and cultured them in serum-free medium as previously described ([Winzeler and Wang, 2013](#)). Human astrocytes were plated on inserts and co-cultured with RGCs for 14–17 days. For quantification of structural synapses, RGCs were fixed and stained with antibodies against the presynaptic marker Bassoon and postsynaptic marker Homer. Synapse number and size were quantified by a custom-written MATLAB program (available upon request). See detailed protocol in [Supplemental Information](#).

#### In Situ Hybridization

Full-length human cDNA expression plasmids were obtained from Dharmicon. Digoxigenin (DIG)-labeled single-stranded antisense riboprobes were prepared, and fresh-frozen 12- $\mu$ m-thick brain sections were processed as previously described ([Zhang et al., 2014](#)) ([Supplemental Information](#)).

#### ACCESSION NUMBERS

We deposited the RNA-seq data in the NCBI GEO database as GEO: GSE73721.

#### SUPPLEMENTAL INFORMATION

Supplemental Information includes Supplemental Experimental Procedures, six figures, and nine tables and can be found with this article online at <http://dx.doi.org/10.1016/j.neuron.2015.11.013>.

#### AUTHOR CONTRIBUTIONS

Y.Z., S.A.S., and B.A.B. conceived the project with input from M.G.H.G. and G.A.G. Y.Z., S.A.S., and B.A.B. designed the experiments and wrote the manuscript. L.E.C. performed the electrophysiology experiments, C.A.P. performed some of the qPCR experiments, Y.Z., S.A.S., and C.C. performed all other experiments. M.G.H.G., G.A.G., E.F.C., L.M.S., S.H.C., J.A.D., G.L., M.S.B.E., G.K.S., H.V., and P.D.B. evaluated patients and resected and examined brain tissue. M.G.H.G. and G.A.G. oversaw tissue procurement.

#### ACKNOWLEDGMENTS

We would like to thank Maria Fabian, Yingmei Li, Annie Abay, Tom Babcock, Andrew Olson, the Stanford University Neurosurgery Brain Bank, and the Stanford Neuroscience Microscopy Service for technical assistance; Won-suk Chung for help with the synaptosome engulfment assay; Mariko Bennett and Chris Bennett for help with *in situ* hybridization; the B.A.B. laboratory for helpful discussions; and Nenad Sestan for brain tissue. This study was supported by: NIH grants R01NS081703 and R01MH099555 and a grant from the JPB Foundation to B.A.B.; a Walter V. and Idun Berry Fellowship and NIH grant K99NS089780 to Y.Z.; and NIH grant T32GM007365, with additional support from the Stanford University School of Medicine and its Medical Scientist Training Program, and NIH grant F30MH106261 to S.A.S. We thank Vincent and Stella Coates and the Dr. Miriam and Sheldon G. Adelson Medical Research Foundation for their generous support.

Received: June 21, 2015

Revised: October 5, 2015

Accepted: November 4, 2015

Published: December 10, 2015

#### REFERENCES

- Agulhon, C., Fiacco, T.A., and McCarthy, K.D. (2010). Hippocampal short- and long-term plasticity are not modulated by astrocyte Ca<sup>2+</sup> signaling. *Science* 327, 1250–1254.
- Allen, N.J., and Barres, B.A. (2009). Neuroscience: Glia - more than just brain glue. *Nature* 457, 675–677.
- Allen, N.J., Bennett, M.L., Foo, L.C., Wang, G.X., Chakraborty, C., Smith, S.J., and Barres, B.A. (2012). Astrocyte glypicans 4 and 6 promote formation of excitatory synapses via GluA1 AMPA receptors. *Nature* 486, 410–414.
- Attwell, D., Buchan, A.M., Chrapak, S., Lauritzen, M., Macvicar, B.A., and Newman, E.A. (2010). Glial and neuronal control of brain blood flow. *Nature* 468, 232–243.
- Banker, G.A. (1980). Trophic interactions between astroglial cells and hippocampal neurons in culture. *Science* 209, 809–810.
- Beach, T.G., and McGeer, E.G. (1988). Lamina-specific arrangement of astrocytic gliosis and senile plaques in Alzheimer's disease visual cortex. *Brain Res.* 463, 357–361.

- Christopherson, K.S., Ullian, E.M., Stokes, C.C.A., Mallowney, C.E., Hell, J.W., Agah, A., Lawler, J., Mosher, D.F., Bornstein, P., and Barres, B.A. (2005). Thrombospondins are astrocyte-secreted proteins that promote CNS synaptogenesis. *Cell* 120, 421–433.
- Chung, W.-S., Clarke, L.E., Wang, G.X., Stafford, B.K., Sher, A., Chakraborty, C., Joung, J., Foo, L.C., Thompson, A., Chen, C., et al. (2013). Astrocytes mediate synapse elimination through MEGF10 and MERTK pathways. *Nature* 504, 394–400.
- Chung, W.-S., Allen, N.J., and Eroglu, C. (2015). Astrocytes control synapse formation, function, and elimination. *Cold Spring Harb. Perspect. Biol.* 7, a020370.
- Clarke, L.E., and Barres, B.A. (2013). Emerging roles of astrocytes in neural circuit development. *Nat. Rev. Neurosci.* 14, 311–321.
- Ding, F., O'Donnell, J., Thrane, A.S., Zeppenfeld, D., Kang, H., Xie, L., Wang, F., and Nedergaard, M. (2013).  $\alpha$ 1-adrenergic receptors mediate coordinated Ca<sup>2+</sup> signaling of cortical astrocytes in awake, behaving mice. *Cell Calcium* 54, 387–394.
- Foo, L.C., Allen, N.J., Bushong, E.A., Ventura, P.B., Chung, W.-S., Zhou, L., Cahoy, J.D., Daneman, R., Zong, H., Ellisman, M.H., and Barres, B.A. (2011). Development of a method for the purification and culture of rodent astrocytes. *Neuron* 71, 799–811.
- Ge, W.-P., Miyawaki, A., Gage, F.H., Jan, Y.N., and Jan, L.Y. (2012). Local generation of glia is a major astrocyte source in postnatal cortex. *Nature* 484, 376–380.
- Gordienko, D.V., and Bolton, T.B. (2002). Crosstalk between ryanodine receptors and IP(3) receptors as a factor shaping spontaneous Ca(2+)-release events in rabbit portal vein myocytes. *J. Physiol.* 542, 743–762.
- Hamilton, N.B., and Attwell, D. (2010). Do astrocytes really exocytose neurotransmitters? *Nat. Rev. Neurosci.* 11, 227–238.
- Han, X., Chen, M., Wang, F., Windrem, M., Wang, S., Shanz, S., Xu, Q., Oberheim, N.A., Bekar, L., Betstadt, S., et al. (2013). Forebrain engraftment by human glial progenitor cells enhances synaptic plasticity and learning in adult mice. *Cell Stem Cell* 12, 342–353.
- Hawrylycz, M.J., Lein, E.S., Guillozet-Bongaarts, A.L., Shen, E.H., Ng, L., Miller, J.A., van de Lagemaat, L.N., Smith, K.A., Ebbert, A., Riley, Z.L., et al. (2012). An anatomically comprehensive atlas of the adult human brain transcriptome. *Nature* 489, 391–399.
- Haydon, P.G., and Nedergaard, M. (2014). How do astrocytes participate in neural plasticity? *Cold Spring Harb. Perspect. Biol.* 7, a020438.
- Huang, W., Sherman, B.T., and Lempicki, R.A. (2009). Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. *Nat. Protoc.* 4, 44–57.
- Huttenlocher, P.R. (1979). Synaptic density in human frontal cortex - developmental changes and effects of aging. *Brain Res.* 163, 195–205.
- Huttenlocher, P.R., de Courten, C., Garey, L.J., and Van der Loos, H. (1982). Synaptogenesis in human visual cortex—evidence for synapse elimination during normal development. *Neurosci. Lett.* 33, 247–252.
- Kang, H.J., Kawasawa, Y.I., Cheng, F., Zhu, Y., Xu, X., Li, M., Sousa, A.M.M., Pletikos, M., Meyer, K.A., Sedmak, G., et al. (2011). Spatio-temporal transcriptome of the human brain. *Nature* 478, 483–489.
- Khakh, B.S., and McCarthy, K.D. (2015). Astrocyte calcium signaling: from observations to functions and the challenges therein. *Cold Spring Harb. Perspect. Biol.* 7, a020404.
- Kola, I., and Landis, J. (2004). Can the pharmaceutical industry reduce attrition rates? *Nat. Rev. Drug Discov.* 3, 711–715.
- Kucukdereli, H., Allen, N.J., Lee, A.T., Feng, A., Ozlu, M.I., Conatser, L.M., Chakraborty, C., Workman, G., Weaver, M., Sage, E.H., et al. (2011). Control of excitatory CNS synaptogenesis by astrocyte-secreted proteins Hevin and SPARC. *Proc. Natl. Acad. Sci. USA* 108, E440–E449.
- Kuffler, S.W., Nicholls, J.G., and Orkand, R.K. (1966). Physiological properties of glial cells in the central nervous system of amphibia. *J. Neurophysiol.* 29, 768–787.
- Malik, N., Wang, X., Shah, S., Efthymiou, A.G., Yan, B., Heman-Ackah, S., Zhan, M., and Rao, M. (2014). Comparison of the gene expression profiles of human fetal cortical astrocytes with pluripotent stem cell derived neural stem cells identifies human astrocyte markers and signaling pathways and transcription factors active in human astrocytes. *PLoS ONE* 9, e96139.
- McCarthy, K.D., and de Vellis, J. (1980). Preparation of separate astroglial and oligodendroglial cell cultures from rat cerebral tissue. *J. Cell Biol.* 85, 890–902.
- Meyer-Franke, A., Kaplan, M.R., Pfrieger, F.W., and Barres, B.A. (1995). Characterization of the signaling interactions that promote the survival and growth of developing retinal ganglion cells in culture. *Neuron* 15, 805–819.
- Molofsky, A.V., Krencik, R., Ullian, E.M., Tsai, H.-H., Deneen, B., Richardson, W.D., Barres, B.A., and Rowitch, D.H. (2012). Astrocytes and disease: a neurodevelopmental perspective. *Genes Dev.* 26, 891–907.
- Mulligan, S.J., and MacVicar, B.A. (2004). Calcium transients in astrocyte end-feet cause cerebrovascular constrictions. *Nature* 431, 195–199.
- Nedergaard, M. (1994). Direct signaling from astrocytes to neurons in cultures of mammalian brain cells. *Science* 263, 1768–1771.
- Nimmerjahn, A., Mukamel, E.A., and Schnitzer, M.J. (2009). Motor behavior activates Bergmann glial networks. *Neuron* 62, 400–412.
- Oberheim, N.A., Wang, X., Goldman, S., and Nedergaard, M. (2006). Astrocytic complexity distinguishes the human brain. *Trends Neurosci.* 29, 547–553.
- Oberheim, N.A., Takano, T., Han, X., He, W., Lin, J.H.C., Wang, F., Xu, Q., Wyatt, J.D., Pilcher, W., Ojemann, J.G., et al. (2009). Uniquely hominid features of adult human astrocytes. *J. Neurosci.* 29, 3276–3287.
- Parpura, V., Basarsky, T.A., Liu, F., Jęftinija, K., Jęftinija, S., and Haydon, P.G. (1994). Glutamate-mediated astrocyte-neuron signalling. *Nature* 369, 744–747.
- Paukert, M., Agarwal, A., Cha, J., Doze, V.A., Kang, J.U., and Bergles, D.E. (2014). Norepinephrine controls astroglial responsiveness to local circuit activity. *Neuron* 82, 1263–1270.
- Pellerin, L., and Magistretti, P.J. (1994). Glutamate uptake into astrocytes stimulates aerobic glycolysis: a mechanism coupling neuronal activity to glucose utilization. *Proc. Natl. Acad. Sci. USA* 91, 10625–10629.
- Rothstein, J.D., Dykes-Hoberg, M., Pardo, C.A., Bristol, L.A., Jin, L., Kuncl, R.W., Kanai, Y., Hediger, M.A., Wang, Y., Schielke, J.P., and Welty, D.F. (1996). Knockout of glutamate transporters reveals a major role for astroglial transport in excitotoxicity and clearance of glutamate. *Neuron* 16, 675–686.
- Schummers, J., Yu, H., and Sur, M. (2008). Tuned responses of astrocytes and their influence on hemodynamic signals in the visual cortex. *Science* 320, 1638–1643.
- Sun, W., McConnell, E., Pare, J.F., Xu, Q., Chen, M., Peng, W., Lovatt, D., Han, X., Smith, Y., and Nedergaard, M. (2013). Glutamate-dependent neuroglial calcium signaling differs between young and adult brain. *Science* 339, 197–200.
- Tien, A.C., Tsai, H.H., Molofsky, A.V., McMahon, M., Foo, L.C., Kaul, A., Dougherty, J.D., Heintz, N., Gutmann, D.H., Barres, B.A., and Rowitch, D.H. (2012). Regulated temporal-spatial astrocyte precursor cell proliferation involves BRAF signalling in mammalian spinal cord. *Development* 139, 2477–2487.
- Ullian, E.M., Sapperstein, S.K., Christopherson, K.S., and Barres, B.A. (2001). Control of synapse number by glia. *Science* 291, 657–661.
- Winzeler, A., and Wang, J.T. (2013). Purification and culture of retinal ganglion cells from rodents. *Cold Spring Harb. Protoc.* 2013, 643–652.
- Zamanian, J.L., Xu, L., Foo, L.C., Nouri, N., Zhou, L., Giffard, R.G., and Barres, B.A. (2012). Genomic analysis of reactive astrogliosis. *J. Neurosci.* 32, 6391–6410.
- Zhang, Y., Chen, K., Sloan, S.A., Bennett, M.L., Scholze, A.R., O'Keefe, S., Phatnani, H.P., Guarnieri, P., Caneda, C., Ruderisch, N., et al. (2014). An RNA-sequencing transcriptome and splicing database of glia, neurons, and vascular cells of the cerebral cortex. *J. Neurosci.* 34, 11929–11947.
- Zonta, M., Angulo, M.C., Gobbo, S., Rosengarten, B., Hossmann, K.-A., Pozzan, T., and Carmignoto, G. (2003). Neuron-to-astrocyte signaling is central to the dynamic control of brain microcirculation. *Nat. Neurosci.* 6, 43–50.